INDEX

Bald, J. G., plan of growth, maturity, and yield of the potato plant, 43–8.
Beans, manuring of, 195–206; annual experiments, 200–3; current manurial practice, 205–7; long-term experiments: Denmark, 199–200; Rothamsted, 196–8; Saxmundham, 198–9.
Bertram, G. C. L., carp farming in Palestine, 187–94.
Blackarm disease, breeding cotton resistant to, 153–74. I. Introductory, 153–60. American Upland cottons, 158–9; blackarm: genetics, 155–8; nature, 154–5; prevalence, 153. II. Breeding methods, 161–74. Adding B_3 by hybridization, 164–6, B_2 and B_3 to a fully susceptible variety, 166–70, B_3 to an Upland variety, 171–2; grading plants for resistance, 162–3; inoculum and spraying, 161–2; selection for B_2 resistance in Upland types, 163–4.
Boyd, D. A., manuring of beans and peas, 195–207.

Carp farming in Palestine, 187–94; carp, 189; fish farming: future of, in Palestine, 193–4, and public health, 188–9, use of land for, 188; growth, yield, harvesting, and management, 192–3; pond construction, 189–90; water-supplies, 191.
Cinchona in Uganda, 75–84; climatic and soil requirements, 78–80; costs, yields, and returns, 83–4; cultivation, 80–2; historical, 75–6; pests and diseases, 82–3; selection, 76–8.
Common, R. H., feeding-value of flax by-products. II. Flax shive, 71–4.
Cotton, *see* Blackarm disease of, 153–74.

Diseases and pests; blackarm of cotton, 153–74; due to night soil, 7; of cinchona in Uganda, 82–3; Solway pine of lambs, 145–52; of Sorghum in Egypt, 211.

Farm school, the Oyo, Nigeria, 135–44; class-room work, 138–9; difficulties, 141–2; finance, 140–1, 142–4; liaison with local farmers, 139; post-school settlement, 139–42; rural rehabilitation, 135; students, 135–9, 140, 141; training farms, 137–8.
Fertilizers and manures; for arable soils in Cyprus, 123 sqq.: for carp ponds, 192; in general, 1 sqq., 104–6, 115–16, 175 sqq.; for Hevea, 182–6; for irrigated soils, 111; for mangolds, 49 sqq.; for peas and beans, 195–207; in perhumid areas, 117–18; for rice, 119–21; for tung trees, 24; for wheat in Middle East, 41.
Flax shive, feeding-value of, 71–4; composition and digestibility, alkali-treated and untreated, 71–3; ryegrass-seed cleanings, 71–2.

Gibberd, A. V., the Oyo farm school, Nigeria, 135–44.

Haines, W. B., manuring Hevea. IV. Conspectus of experimental improvements achieved in mature stands at the end of ten years, with a special note on seed production, 182–6.
Hevea, manuring of. IV. Improvements achieved in ten years, 182–6; benefits, general, 182–3, measured, 183–4, 185–6; seed-production, 185–6.

Knight, R. L., breeding cotton resistant to blackarm disease. I. Introductory, 153–60. II. Breeding methods, 161–74.

Littlejohn, L., some aspects of soil fertility in Cyprus, 123–34.

Mangolds, Rothamsted experiments on, 1876–1940. IV. Composition of mangolds grown on Barnfield, 49–70. (i) Dry-matter content of leaves and roots, 49–56; effects of F.Y.M. and fertilizers, 49–50; seasonal variation, 50–1; sugar-content, 53–6; water-content in growing-season, 53. (ii) Nitrogen-content of leaves and roots, 57–70; effects of F.Y.M. and fertilizers, 57–60, 70; mineral-content, 65–70; recovery of added nitrogen, 60–2; seasonal variation, 62–5.
Mann, H. H., wheat in the Middle East, 31–42.
— millets in the Middle East, 208–16.
Millets in the Middle East, 208–16; advantages over other cereals, 208; future of, 215; origins of, 208; Panicum millets, 212–15; *Sorghum vulgare*, 209–12.
Mitchell, R. L., *see* Stewart, J.
Munro, S. S., relative influence of heredity and environment on fertility and hatchability in Wyandottes, 25–30.

Peas, fertilizer requirements of, 203–4; current manurial practice, 206.
Plank, J. E. van der, some climatic factors determining high yields of potatoes, 217–23.
Potato plant, the, a plan of growth, maturity, and yield, 43–8; growth, 42–6, 47–8; maturity, 46; yield, 46–7.
Potatoes, climatic factors determining high yields of. I. Temperature and length of growing-season, 217–23; in maritime and continental climates at high latitudes, 218–20; possibility of improving yields in high-latitude continental climates by breeding, 220–3.

Rainfall of India, 85–99; periodicity, 95; physical and climatic features, 85–8; regional peculiarities, 97–9; SW. monsoon, 88–95, 96; unusually heavy rainfall, 95, 97; utilization of, 99.

Ramdas, L. A., rainfall of India: a brief review, 85–99.
Richardson, H. L., soil-fertility maintenance under different systems of agriculture, 1–17.
— improvement of soil fertility. I. General considerations, 100–8. II. Relationships with climate, 109–22. III. Agriculture, food, and health, 175–80.

Soil fertility, improvement of. I. General considerations, 100–8; crop production, goals and limits, 101–2; manuring, 104–6; reclamation of waste lands, 102–3; soil conservation, 106–7; survey and investigation, 103–4. II. Relationships with climate, 109–22; rainfall areas, very low, 109–11, moderately low, 111–14, moderate, 114–17, heavy, 117–21. III. Agriculture, food, and health, 175–81; manures, organic and inorganic, 175–7; manuring, health and disease, 177–8; quantity and quality of produce, 175–7; conclusions to Pts. I, II.
— III, maintenance of, 1–17; fertility of natural soils, 3; fuel problem, 14; grassland farming, 12–13; modern agriculture, 8–11; natural recuperation, 2; plantation agriculture, 12; primitive agriculture, 3–4; relation of fertility to relief, 14; rice, 11–12; signs of low fertility, 2–3; soil erosion and conservation, 13–14; traditional agriculture, 4–8; China, 6–8; Egypt, 4–5; Europe, 5–6; India, 6; wheat yields, 5, 6, 8, 9, 10, 11, 15–16.
Soils, fertility of arable, in Cyprus, 123–34; fallowing, 127–8, 131; green manuring, 131, 132–3; manurial trials with a compound fertilizer, farmyard and town refuse, sheep manure, 129–30; on wheat, 123–7; organic manuring, efficiency of, 128 sqq.
Solway pine of lambs in Kircudbrightshire, 145–52; analyses of soils and pastures, 145–7; cobalt pining areas of Scotland, 151, 152; trials with cobalt dosing, 147–8; cobalt manuring, 148–51.
Stewart, A. B., *see* Stewart, J.
Stewart, J. (with Mitchell, R. L., Stewart, A. B., and Young, H. M.), Solway pine: a marasmic condition in lambs in certain districts of Kirkcudbrightshire, 145–52.

Thomas, A. S., cinchona in Uganda, 75–84.
Tung tree, cultivation of, in Nyasaland, 18–24; cultivation, 23–4; manuring, 24; nursery work, 22–3; pollination, 24; production of improved planting material, 19–22; seedling plantations, 18–19; spacing, 23; transplanting, 23.

Watson, D. J., Rothamsted experiments on mangolds, 1876–1940. IV. Composition of the mangolds grown on Barnfield. (i) Dry-matter content of leaves and roots, 49–56. (ii) N-content of leaves and roots, 57–70.
Webster, C. C., cultivation of the tung tree in Nyasaland, 18–24.
Wheat in the Middle East, 31–42; conditions in Abyssinia and Eritrea, 40; Afghanistan, 39–40; Egypt, 40; Iran, 39; Iraq, 39; Palestine, 37–8; Syria and Lebanon, 37; Transjordan, 38–9; Turkey, 36–7; influence of climate, 32–3; production, 33–5; technical difficulties, 35–6; varieties, 31–2.
Wyandottes, white, fertility and hatchability of, 25–30; experimental, 27–9; farm records, 25–7; relation of fertility and hatchability, 29.

Young, H., *see* Stewart, J.

CORRIGENDUM.

Volume XIV, No 53, line 8 from bottom *in lieu of* '13/5', *read* '5/13'.